

COVID-19, social distance and adolescents' risk behaviours, wellbeing and life satisfaction: a proxy study drawn from HBSC study

COVID-19, distancia social y conductas de riesgo de los adolescentes, bienestar y satisfacción con la vida: un estudio proxy extraído del estudio HBSC

Gina Tomé, PhD

FMH/Universidade de Lisboa, ISAMB/Universidade de Lisboa, Portugal. SFRH/BPD/108637/2015

Cátia Branquinho, PhD

Ana Cerqueira, Msc

Margarida Gaspar de Matos, PhD

FMH/Universidade de Lisboa, ISAMB/Universidade de Lisboa, Portugal

RESUMEN

Privar a las personas de su libertad tiene efectos devastadores sobre el bienestar y la salud mental, especialmente en los adolescentes. Esta fue la situación con la reciente pandemia de COVID-19 que obligó a los adolescentes a quedarse en casa.

Para simular una situación de ausencia de interacciones sociales fuera del contexto familiar, se utilizaron datos portugueses del estudio HBSC / WHO 2018. Se pretendía explorar y comprender cuál de los factores de riesgo y de protección de la salud de los adolescentes habituales se vería más afectado entre aquellos que no tienen contacto con sus compañeros después de la escuela.

Los resultados muestran que, por un lado, el "distanciamiento social de los compañeros" en general reduce los riesgos para la salud, como el consumo de refrescos, el consumo de alcohol, tabaco y drogas y la participación en la violencia (peleas, victimización por bullying y lesiones). Por otro lado, disminuye la percepción de bienestar y satisfacción con la vida y, en general, aumenta los síntomas psicológicos.

PALABRAS CLAVE

Distanciamiento social; conductas de riesgo; conductas protectoras; bienestar; adolescencia.

ABSTRACT

Depriving people of their liberty has devastating effects upon wellbeing and mental health, especially in adolescents. This was the situation with the recent COVID-19 pandemic that forced adolescents to stay at home.

In order to simulate a situation of absence of social interactions outside the family context, Portuguese data from the HBSC / WHO 2018 study were used. It was intended to explore and understand which of the usual adolescents' health risk and protective factors would be more affected among those who do not have contact with peers after school.

The results show that, on the one hand, "social distancing from colleagues" in general reduces health risks, such as consumption of soft drinks, alcohol, tobacco and drug use and involvement in violence (fights, victimization by bullying and injuries). On the other hand, it decreases the perception of well-being and life satisfaction and in general increases the psychological symptoms.

KEYWORDS

Social distancing; risk behaviours; protective behaviours; wellbeing; adolescence.

Recibido: 20/02/2020; aceptado: 24/08/2020

Correspondencia: Gina Tomé, Faculdade de Motricidade Humana, Universidade de Lisboa, Estrada da Costa
ginatome@sapo.pt

Introduction

Well-being is identified as one of the essential elements for the quality of life of young people (Gaspar et al., 2012; Jiménez-Iglesias, Camacho, Rivera, Moreno & Matos, 2017). Depriving people of their liberty has potential devastating effects upon wellbeing and mental health, especially when it comes to adolescents. A very recent study where about 600 adolescents living under lockdown were heard about their experiences (Branquinho, Kelly, Arevalo, Santos, & Matos, 2020, submitted), showed that these adolescents' report mix experiences. surfacing the fact that a few indicators of physical health indeed got better (better nutrition, less substance use and less involvement in violence and accidents) while the general perception of low wellbeing and life satisfaction and a few psychological symptoms, sleep problems and sedentary behaviour (screen time) increased as well.

Providing relevant leisure time occupations and quality schooling is also extremely relevant in childhood and adolescence due to the specific developmental issues of this phase (Brooks et al., 2020), and this issue become extremely difficult during lockdown.

Youth psychosocial development is influenced by individual and environmental/contextual factors associated with the adolescents' well-being, of which school involvement and success are examples (Berger, Alcalay, Torretti, & Milicic, 2011; Lewis, Huebner, Malone, & Valois, 2011).

During the recent COVID-19 pandemic and following the social distancing protective measures, many schools have been closed

and classes moved to e-learning home-based models. This raised the question of the potential associations between the social distancing measures and the children's and adolescents' well-being (Golberstein et al., 2020), because there is an established influence of their life (family, peers and school) and well-being (Choi, 2018; Gaspar, Cerqueira, Branquinho, & Matos, 2018).

Developing a good relationship with the peer group is an essential factor in promoting the youths' emotional well-being (Camacho et al., 2017; Tomé et al., 2018). Young people who have more positive school experiences with greater peer support, report better psychosocial adjustment, and tend to perceive a higher life satisfaction (Marques, Lopez, Fontaine, Coimbra, & Mitchell, 2015; Matos et al., 2017).

Social context is an extremely feature in adolescents' positive development and the negative consequences of lockdown for adolescents' wellbeing, while not yet totally estimated can easily be hypothesized and foreseen from available and tested theoretical models and empirical (Matos et al., 2020, in press; Tomé, et al., 2020; Jensen et al. 2020, submitted).

It is expected that Negative consequences of lockdown for adolescents' physical and mental health and well-being are expected may include involvement in risk behaviours such as increased drug and alcohol use, violence, and poor school performance (Simões, Rivera, Moreno & Matos, 2018, Branquinho et al., 2020, submitted; IREFREA, 2020). This same feature was found in a paper comparing data from HBSC in 2010 and in 2014 (Matos et al., 2012; Matos, Simões, Camacho, Reis & Aventura So-

cial Team, 2015), before and after the impact of the economic recession experienced between 2008 and 2014 (Matos et al., 2015). This paper established that wellbeing and mental health were the first to deteriorate even when there were no signs of physical health problems.

In the context of social distancing due to the recent COVID-19 pandemic, we revisited the 2018 wave of HBSC (Matos & Equipa Aventura Social, 2018) in Portugal, in order to explore the effect of the lack of socialization with friends after school upon several risk and protective factors. We also intend to explore the impact of these risk and protective factors upon wellbeing and life satisfaction, controlling for social contact with peers after school.

Method

This survey is part of the Health Behaviour in School-aged Children (HBSC) study (Currie et al., 2004; Matos et al., 2006; Matos & Aventura Social Team, 2018). The HBSC is a collaborative WHO study, undertaken in 44 countries and that aims to study the school-aged children behaviour regarding health and risk behaviours. Portugal has been part of this group of countries since 1996. The HBSC is a school-based survey of adolescents' health behaviours, carried out every 4 years. Collected data is used at a national and international level, using an internationally standardized methodological protocol (Roberts et al., 2007) that intends to: (1) gain a new vision into young people's health and well-being, (2) understand the social and psychological determinants of health and (3) incorporate policies to improve young people's lives.

Participants

The 2018 study provided national representative data of 8215 Portuguese adolescents, randomly chosen from those attending 6th grade, 8th grade (middle school), 10th grade and 12th grade (high school) during the 2017/2018 academic year. The sample included 52.7% of girls and 47.3% of boys, whose mean age was 14.36 years old (standard deviation 2.28). This study used a subset of 8th (n=2766), 10th (n=1711) and 12th graders (n=1218) to represent middle school and high school educational stages. They were randomly selected from 42 national vertical clusters of schools, in a total of 476 classes, in a national sample geographically stratified by Education Regional Divisions in Portugal. The overall procedure has been described elsewhere (Currie et al., 2004; Matos et al., 2012); in brief, this study has the approval of a scientific committee, an ethical national committee and the national commission for data protection and followed strictly all the guidelines for protection of human rights; adolescents' participation in the survey and completion of the questionnaires was voluntary and anonymous. The sample was nationally representative of the respective grade levels.

Variables and Measures

For this study, variables that evaluate the mental health and the well-being of Portuguese adolescents were used: Well-being was measured by the Kidscreen scale, the global Well-being with 10 items scale with answer options: 1-Never; 2-Rarely; 3-Quite often; 4-Very often; 5-Always; Higher score higher wellbeing

(Gaspar et al., 2012). Life satisfaction was evaluated with the Cantril scale (1965), graphically represented as a ladder, where step '10' corresponds to "best possible life" and step '0' represents "the worst possible". Psychological symptoms were assessed with the questions used in the HBSC study protocol for health symptoms, were higher score higher symptoms (sadness, irritation or bad temper, nervousness, tiredness and exhaustion, difficulties in getting to sleep and self-harm) (Currie et al., 2004; Matos et al., 2012). Variables associated with health risks were: drinking coke or other soft drinks that contains sugar; alcohol, tobacco and drugs consumption, injuries, fights involvement, being bullied and being cyberbullied, were assessed with the questions used in the HBSC study protocol (Currie et al., 2004; Matos et al., 2012) described in table 1. The detailed description of the variables used is shown in table 1.

Data analysis

The data was analyzed using the Statistical Package for Social Sciences (SPSS) version 24 for Windows. Descriptive correlational and comparative analyzes were performed (ANOVAS and Chi-Square) and finally multiple linear and logistic regression models.

Results

To analyze the differences between adolescents who are more or less often with friends, three groups were formed for the variables "peer contact frequency-after school: None or 1 day; A few days; Every day.

For the differences between frequency of after school contact with the peers, it was found by Chi-Square that adolescents who spend less days with friends after school (none or 1 day), consume less: coke or other soft drinks that contain sugar (never or less than once a week) ($\chi^2=91.688(4)$, $p\leq.001$, 49.4%), alcohol – in the last 30 days (never/1-2 days) ($\chi^2=102.975(4)$, $p\leq.001$, 90.8%), tobacco – in the last 30 days (never/1-2 days) ($\chi^2=98.828(4)$, $p\leq.001$, 95.7%), marijuana – in the last 30 days (never/1-2 days) ($\chi^2=35.638(4)$, $p\leq.001$, 98%), had less injuries (never/1 time) ($\chi^2=42.111(4)$, $p\leq.001$, 84.7%), involved less in fights (never/1 time) ($\chi^2=26.441(4)$, $p\leq.001$, 91.4%), had more feelings: sadness (about every day) ($\chi^2=44.495(4)$, $p\leq.001$, 15.1%), irritability or bad temper (about every day) ($\chi^2=24.831(4)$, $p\leq.001$, 17.4%), nervousness (about every day) ($\chi^2=31.824(4)$, $p\leq.001$, 20.4%), difficulties in getting to sleep (about every day) ($\chi^2=9.522(4)$, $p\leq.05$, 14.4%), finally, are less satisfied with life (nothing satisfied) ($\chi^2=44.216(4)$, $p\leq.001$, 32.8%) and mention less quality of life – Kidscreen (poor quality) ($\chi^2=95.860(4)$, $p\leq.001$, 41%). Finally, adolescents who spend more days with friends after school (every day), are more involved in bullying and cyberbullying behaviour: being bullied (several times a week) ($\chi^2=22.617(4)$, $p\leq.001$, 4.4%) and being cyberbullied (several times a week) ($\chi^2=9.632(4)$, $p\leq.05$, 1.3%) and feel more often tiredness and exhaustion (about every day) ($\chi^2=9.632(4)$, $p\leq.01$, 23.3%).

Table 3 shows the differences between the peer contact frequency – after school, for life satisfaction and Kidscreen. Regarding life

Table 1
Variables used in the analysis

Item	Original Response Options	Recorded Response Options
Kidscreen (Global well-being)	Global Well-being 10 items scale	Higher score higher well-being
Life Satisfaction	1-Never; 2-Rarely; 3-Quite often; 4-Very often; 5-Always; Step '10' corresponds to "best possible life and step '0' represents "the worst possible life".	Higher score higher life satisfaction
Peer contact frequency (after school)	Cantril scale, graphically represented as a ladder How many days a week do you usually spend time with friends right after school? 1- 0 days; 2- 1; 3- 2; 4- 3; 5- 4; 6 - 5; 7- 6 days.	1 - none or 1 day; 2 - a few days; 3- every day; /or dichotomized variable for regression analysis; 0- 0 days; 1- a few days;
Coke or other soft drinks that contain sugar (transformed into Zscore for regression)	How many times a week do you usually eat or drink? 1- Never; 2- Less than once a week; 3- Once a week; 4- 2-4 days a week; 5-5-6 days a week; 6- Once a day, every day; 7- Every day, more than once.	Higher score higher contact Higher score higher consumption
Alcohol - in the last 30 days (transformed into Zscore for regression)	How many days there was (if any) that you have drunk alcohol: 1- Never; 2- 1-2 days; 3- 3-5 days; 6-9 days; 10-19 days; 20-29 days; 30 days (or more)	Higher score higher alcohol abuse
Tobacco - in the last 30 days (transformed into Zscore for regression)	How many days there was (if any) that you have smoke: 1- Never; 2- 1-2 days; 3- 3-5 days; 6-9 days; 10-19 days; 20-29 days; 30 days (or more)	Higher score higher consumption

(Continue)

Table 1
Variables used in the analysis (Continuation)

Drugs – in the last 30 days (transformed into Zscore for regression)	How many days there was (if any) that you have consumed "marijuana":	1- Never; 2- 1-2 days; 3- 3-5 days; 6-9 days; 10-19 days; 20-29 days; 30 days (or more)	Higher score higher consumption
Injuries (transformed into Zscore for regression)	During the past 12 months, how many times were you injured and had to be treated by a doctor or nurse?	1- I was not injured in the past 12 months; 2-1 time; 3- 2 times; 4-3 times; 5- 4 times or more	Higher score higher injuries frequency
Fights (transformed into Zscore for regression)	During the past 12 months, how many times were you in a physical fight?	1- I have not been in a physical fight in the past 12 months; 2-1 time; 3-2 times; 4-3 times; 5-4 times or more.	Higher score higher fights frequency
Being bullied (transformed into Zscore for regression)	How often have you been bullied at school in the past couple of months?	1- I have not been bullied at school in the past couple of months; 2-It has only happened once or twice; 3-2 or 3 times a month; 4-About once a week; 5-Several times a week;	Higher score higher bullying-victimization frequency
Cyberbullying victim (transformed into Zscore for regression)	How often have you been cyberbullied at school in the past couple of months?	1- I have not been cyberbullied at school in the past couple of months; 2-It has only happened once or twice; 3-2 or 3 times a month; 4-About once a week; 5-Several times a week;	Higher score higher cyberbullying-victimization frequency
Sadness (transformed into Zscore for regression)	In the last 6 months: how often have you had the following....?	1-About every day; 2-More than once a week; 3-About every week; 4-About every month; 5-Rarely or never.	1- Rarely or never; 2- About every month; 3- About every week; 4- More than once a week; 5- About every day. Higher score higher sadness
Irritability or bad temper (transformed into Zscore for regression)	In the last 6 months: how often have you had the following....?	1-About every day; 2-More than once a week; 3-About every week; 4-About every month; 5-Rarely or never.	1- Rarely or never; 2- About every month; 3- About every week; 4- More than once a week; 5- About every day. Higher score higher bad temper

(Continue)

Table 1
Variables used in the analysis (Continuation)

Nervousness (transformed into Zscore for regression)	In the last 6 months: how often have you had the following....?	1-About every day; 2-More than once a week; 3-About every week; 4-About every month; 5-Rarely or never.	1- Rarely or never; 2- About every month; 3- About every week; 4- More than once a week; 5- About every day. Higher score higher nervousness
Tiredness and exhaustion (transformed into Zscore for regression)	In the last 6 months: how often have you had the following....?	1-About every day; 2-More than once a week; 3-About every week; 4-About every month; 5-Rarely or never.	1- Rarely or never; 2- About every month; 3- About every week; 4- More than once a week; 5- About every day. Higher score higher tiredness
Difficulties in getting to sleep (transformed into Zscore for regression)	In the last 6 months: how often have you had the following....?	1-About every day; 2-More than once a week; 3-About every week; 4-About every month; 5-Rarely or never.	1- Rarely or never; 2- About every month; 3- About every week; 4- More than once a week; 5- About every day. Higher score higher difficulties getting to sleep

satisfaction, it can be seen that adolescents who were most satisfied with life spend more time with friends after school (every day) ($M=7.5$, $SD=1.9$), $F(2,4502) = 19.793$, $p < .001$. As for Kidscreen, the same pattern can be seen, meaning that adolescents with a better perception of quality of life spend more time with friends after school (every day) ($M=37.6$, $SD=7.3$), $F(2,4502) = 48.298$, $p < .001$.

To understand the predictive effect of the variables used in this study regarding life satisfaction and quality of life (Kidscreen), multiple linear regression analysis was conducted, including variables that were significant at a bivariate level (Chi-Square and ANOVA).

The regression equation for the model of the life satisfaction explained 21% of the variance ($R^2=.215$). In this model, the explanation of the life satisfaction was obtained through the peer contact frequency – after school (stay with friends after school) ($\beta=.038$, $p=.005$), drinking coke or other soft drinks that contain sugar (higher consumption) ($\beta=.034$, $p=.012$), tobacco – in the last 30 days (lower consumption) ($\beta= -.055$, $p=.001$), alcohol – in the last 30 days (lower consumption) ($\beta= -.037$, $p=.015$), being bullied (less involvement) ($\beta= -.079$, $p<.001$), have feelings of: sadness (less feelings) ($\beta= -.314$, $p<.001$), irritability or bad temper (less feelings) ($\beta= -.093$, $p<.001$), tiredness and exhaustion (less feelings) ($\beta= -.050$, $p=.002$) and difficulties in getting to sleep (less feelings) ($\beta= -.077$, $p<.001$).

The regression equation for the model of the quality of life (Kidscreen) explained 27% of the variance ($R^2=.270$). In this model, the explanation of the quality of life (Kidscreen) was

Table 2
Distributions between groups of After School - Peer contact frequency Qui-squares

	Peer contact frequency (after school)						Total	X ²	df.	
	None or 1 day		A few days		Every day					
	N	%	N	%	N	%				
Coke or other soft drinks that contain sugar	Never or less than once a week	690	49.4	980	39.6	193	30.3	1663	91.688***	4
	Few days	518	37.1	1117	45.2	288	45.3	1923		
	Every day	188	13.5	376	15.2	155	24.4	719		
Alcohol – in the last 30 days	Never/1-2 days	1268	90.8	2069	83.7	496	78	3833	102.975***	4
	Few days	97	6.9	350	14.2	93	14.6	540		
	Many days	31	2.2	54	2.2	47	7.4	132		
Tobacco – in the last 30 days	Never/1-2 days	1336	95.7	2288	92.5	541	85.1	4165	98.828***	4
	Few days	29	2.1	95	3.8	25	3.9	149		
	Many days	31	2.2	90	3.6	70	11	191		
Drugs – in the last 30 days	Never/1-2 days	1368	98	2404	97.2	595	93.6	4367	35.638***	4
	Few days	13	0.9	49	2	23	3.6	85		
	Many days	15	1.1	20	0.8	18	2.8	53		
Injuries	Never/1 time	1182	84.7	1938	78.4	484	76.1	3604	42.111***	4
	Sometimes	151	10.8	421	17	101	15.9	673		
	4 times or more	63	4.5	114	4.6	51	8	228		
Fights	Never/1 time	1276	91.4	2223	89.9	538	84.6	4037	26.441***	4
	Sometimes	79	5.7	173	7	58	9.1	310		
	4 times or more	41	2.9	77	3.1	40	6.3	158		
Bullying Victim	I have not been bullied/once or twice	1293	92.6	2358	95.3	591	92.9	4242	22.617***	4
	Sometimes	60	4.3	70	2.8	17	2.7	147		
	Several times a week	43	3.1	45	1.8	28	4.4	116		
Cyberbullying victim	I have not been cyberbullied/once or twice	1356	97.1	2429	98.2	618	97.2	4403	9.632*	4
	Sometimes	30	2.1	34	1.4	10	1.6	74		
	Several times a week	10	0.7	10	0.4	8	1.3	28		

(Continue)

Table 2
Distributions between groups of After School - Peer contact frequency Qui-squares (Continuation)

Sadness	Rarely/few times	868	62.2	1687	68.2	428	67.3	2983	44.495***	4
	Sometimes	317	22.7	577	23.3	128	20.1	1022		
	About every day	211	15.1	209	8.5	80	12.6	500		
Irritability or bad temper	Rarely/few times	740	53	1379	55.8	366	57.5	2485	24.831***	4
	Sometimes	413	29.6	780	31.5	161	25.3	1354		
	About every day	243	17.4	314	12.7	109	17.1	666		
Nervousness	Rarely/few times	676	48.4	1249	50.5	346	54.4	2271	31.824***	4
	Sometimes	435	31.2	873	35.3	188	29.6	1496		
	About every day	285	20.4	351	14.2	102	16	738		
Tiredness and exhaustion	Rarely/few times	604	43.3	1131	45.7	299	47	2034	19.732**	4
	Sometimes	481	34.5	896	36.2	189	29.7	1566		
	About every day	311	22.3	446	18	148	23.3	905		
Difficulties in getting to sleep	Rarely/few times	909	65.1	1664	67.3	443	69.7	3016	9.522*	4
	Sometimes	286	20.5	503	20.3	103	16.2	892		
	About every day	201	14.4	306	12.4	90	14.2	597		

Table 3
Differences for life satisfaction and kidscreen for After School - Peer contact frequency- ANOVAS

After School -Peer contact frequency	None or 1 day			A few days			Every day			p	
	N	M	SD	N	M	SD	N	M	SD		F
Life satisfaction	1396	7.1	1.9	2473	7.4	1.6	636	7.5	1.9	19.793	.000
Kidscreen	1396	34.9	7.7	2473	37	6.7	636	37.6	7.3	48.298	.000

Table 4

Multiple Linear Regression - Enter Method - Life satisfaction predictors (r2 = 21%)

	Variable included	β	t	p	R ² _a	F(model fit)*
Life satisfaction	Peer contact frequency - After school (dichotomic: 0- 0 days; 1- a few days)	.038	2.811	.005	.215	88.944***
	Coke or other soft drinks that contain sugar (Zscore)	.034	2.504	.012		
	Tobacco – in the last 30 days (Zscore)	-.055	-3.333	.001		
	Alcohol – in the last 30 days (Zscore)	-.037	-2.445	.015		
	Marijuana – in the last 30 days (Zscore)	.017	1.050	.290		
	Being Bullied (Zscore)	-.079	-5.538	.000		
	Cyberbullying victim (Zscore)	-.007	-.527	.598		
	Fights (Zscore)	-.024	-1.695	.090		
	Injuries (Zscore)	.018	1.296	.195		
	Sadness (Zscore)	-.314	-18.167	.000		
	Irritability or bad temper (Zscore)	-.093	-5.231	.000		
	Nervousness (Zscore)	.029	1.687	.092		
	Tiredness and exhaustion (Zscore)	-.050	-3.169	.002		
	Difficulties in getting to sleep (Zscore)	-.077	-5.145	.000		

Table 5

Multiple Linear Regression - Enter Method - Kidscreen predictors (r2= 27%)

	Variable included	β	t	p	R ² _a	F(model fit)*
Kidscreen	Peer contact frequency - After school (dichotomic: 0- 0 days; 1- a few days)	.095	7.343	.000	.270	119.617***
	Coke or other soft drinks that contain sugar (Zscore)	-.002	-.184	.854		
	Tobacco – in the last 30 days (Zscore)	-.021	-1.336	.182		
	Alcohol – in the last 30 days (Zscore)	-.024	-1.637	.102		
	Marijuana – in the last 30 days (Zscore)	-.041	-2.716	.007		
	Being Bullied (Zscore)	-.055	-4.031	.000		
	Cyberbullying victim (Zscore)	-.085	-6.193	.000		
	Fights (Zscore)	-.006	-.421	.674		
	Injuries (Zscore)	.018	1.376	.168		
	Sadness (Zscore)	-.284	-17.051	.000		
	Irritability or bad temper (Zscore)	-.083	-4.831	.000		
	Nervousness (Zscore)	-.048	-2.924	.003		
	Tiredness and exhaustion (Zscore)	-.059	-3.860	.000		
	Difficulties in getting to sleep (Zscore)	-.109	-7.523	.000		

Table 6
 Logistic Regression – After School (YES- stay with friends, at least 1 day after school) - Peer contact frequency Predictors

	β	E.P	Sig	OR	95%IC than	95% IC to
Kidscreen	.044	.006	.000	1.045	1.032	1.059
Life satisfaction	.022	.026	.393	1.022	.972	1.075
Coke or other soft drinks that contain sugar (Zscore)	.141	.042	.001	1.152	1.060	1.250
Tobacco – in the last 30 days (Zscore)	.130	.062	.035	1.139	1.009	1.286
Alcohol – in the last 30 days (Zscore)	.289	.058	.000	1.335	1.191	1.496
Marijuana – in the last 30 days (Zscore)	-.067	.055	.222	.935	.840	1.041
Being Bullied (Zscore)	-.108	.040	.000	.898	.830	.972
Cyberbullying victim (Zscore)	-.030	.043	.489	.971	.892	1.056
Fights (Zscore)	-.021	.045	.639	.979	.896	1.070
Injuries (Zscore)	.143	.045	.002	1.154	1.056	1.261
Sadness (Zscore)	-.058	.054	.281	.944	.849	1.049
Irritability or bad temper (Zscore)	.003	.054	.956	1.003	.901	1.116
Nervousness (Zscore)	-.050	.052	.340	.952	.859	1.054
Tiredness and exhaustion (Zscore)	.004	.049	.932	1.004	.913	1.105
Difficulties in getting to sleep (Zscore)	.089	.046	.052	1.093	.999	1.195
Constant	-.114	.260	.662	.893		

$R^2_{\text{N}} = .06$
 $\chi^2_{\text{HL}} p = 20.795; .008$

obtained through the peer contact frequency – after school (stay with friends after school) ($\beta=.095$, $p<.001$), consumption of marijuana – in the last 30 days (lower consumption) ($\beta= -.041$, $p=.007$), being bullied (less involvement) ($\beta= -.055$, $p<.001$), cyberbullying victim (less involvement) ($\beta= -.085$, $p=.000$), have feelings of: sadness (less feelings) ($\beta= -.284$, $p<.001$), irritability or bad temper (less feelings) ($\beta= -.083$, $p<.001$), nervousness (less feelings) ($\beta= -.048$, $p=.003$), tiredness and exhaustion (less feelings) ($\beta= -.059$, $p<.001$) and difficulties in getting to sleep (less feelings) ($\beta= -.109$, $p<.001$).

Finally, a logistic regression model was performed in order to explain peer contact frequency – after school (stay with friends after school), an adjusted model was obtained (Hosmer & Lemeshow $\chi^2=20.795(8)$, $p=.008$) and the regression equation explained 6% of the variance (Nagelkerke $R^2=.06$). In this model the explanation of the “stay with friends after school” was made by the variables: quality of life (Kidscreen) (better quality) ($\beta=.044$, $p<.001$), drinking coke or other soft drinks that contain sugar (higher consumption) ($\beta=.141$, $p=.001$), tobacco – in the last 30 days (higher consumption) ($\beta=.130$, $p=.035$), alcohol – in the last 30 days (higher consumption) ($\beta=.289$, $p<.001$), being bullied (less involvement) ($\beta= -.108$, $p<.001$) and injuries (more injuries) ($\beta=.143$, $p=.002$).

Discussion

In the context of social distancing due to the recent COVID-19 pandemic, we studied the effect of the lack of socialization with friends

after school upon several risk and protective factors. We also explore the impact of these risk and protective factors upon wellbeing and life satisfaction.

As Branquinho and colleagues (2020, submitted) refer, in a context of lockdown [or confinement] due to the COVID-19 pandemic, adolescents who do not see their peers have lower levels of wellbeing, mental health and life satisfaction. This happens even if they report less risk behavior (drinking sweet drinks and alcohol, smoking tobacco or marijuana) and less involvement in situations of violence and accidents.

Depriving adolescents from the contact with peers and peer social support harms their mental health, and as schools move back to normal and lockdown ends, it is important to provide relevant leisure time occupations with peers, together with quality schooling changed to fit the situation (Brooks et al., 2020).

There is an established influence of the various contexts in which adolescents are involved (family, peers and school) and wellbeing (Choi, 2018; Gaspar, Cerqueira, Branquinho, & Matos, 2018). With the COVID-19 pandemic different patterns were shaped meaning for the great majority of adolescents a better family monitoring but a lower peer contact, which, was checked in the present study, low health risks but also decrease wellbeing, being established that developing a good relationship with the peer group is an essential factor in promoting the youths' emotional well-being (Camacho et al., 2017; Tomé et al., 2018).

With the lockdown any positive school experience was remote, peer support decrease

and as reported previously (Marques, Lopez, Fontaine, Coimbra & Mitchell, 2015; Matos et al., 2017), adolescents report lower psychosocial adjustment, and tend to perceive poorer life satisfaction.

Results show that, on the one hand, “social distancing from peers” in general reduce health risks such as soda consumption, alcohol, tobacco and drug use and violence involvement (fights, bullying victimization and injuries). On the other hand, decreases the perception of wellbeing and life satisfaction and in general, increases the psychological symptoms. This fact has important messages for health and educational professionals, as well as for public policies.

If we can expect an improvement in a general pattern of less health risks with the lockdown, it is also the case that we can expect mental health to deteriorate. Thus, there must be a focus of public policies and interventions in the education and the health sector, when classes return to “normal” in the next academic year. It is advisable to have a period for recovering from the delays related to the academic setbacks. And it is also extremely advisable to keep the psychological wellbeing in mind and to consider a period in which adolescents can be heard with regard to their lockdown experiences, before coming back to their “usual” academic routines.

References

Berger, C., Alcalay, L., Torretti, A., & Milicic, N. (2011). Socio-emotional well-being and academic achievement: Evidence from a multilevel approach. *Psicologia: reflexão e crítica*, 24(2), 344-351.

- Branquinho, C., Kelly, C., Arevalo, L., Santos, A., & Matos, M. G. (2020). “Hey, we also have something to say”: a qualitative study of Portuguese adolescents’ and young people’s experiences under COVID-19. *Journal of Community Health*, 48, 2740-2752. <https://doi.org/10.1002/jcop.22453>.
- Brooks, S., Webster, R., Smith, L., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. (2020). The psychological impact of quarantine and how to reduce it: rapid review of the evidence *Lancet*, 395, 912-20.
- Camacho, I., Matos, M. G., Tomé, G., Reis, M., Jiménez-Iglesias, A., Moreno, M.C., & Galvão, D. (2017). Brothers and Sisters - “More” is Better? Effects on School, Violence and Health. *World Journal of Educational Research*, 4(2), 313-326. Doi: 10.22158/wjer.v4n2p313
- Cantril, H. (1965). *The pattern of human concerns*. New Brunswick, NJ: Rutgers University Press.
- Carvalho, M., Branquinho, C. & Matos, M. G. (2020). Cyberbullying and Bullying: Impact on psychological symptoms and well-being. *Child Indicators Research*, 13. <https://doi.org/10.1007/s12187-020-09756-2>
- Choi, A. (2018). Emotional well-being of children and adolescents: Recent trends and relevant factors. OECD Education Working Papers. Paris: OECD Publishing.
- Currie, C., Roberts, C., Morgan, A., Smith, R., Settertulte, W., Samdal, O., & Rasmussen, V. (2004). HBSC, and WHO cross national study: research protocol for the 2001/2002 survey. Copenhagen, DK: WHO.
- Gaspar, T., Cerqueira, A., Branquinho, C. & Matos, M. G. (2018). The effect of a social-emotional school-based intervention upon social and personal skills in children and adolescents. *Journal of Education and Learning*, 7(6), 57-66. Doi: 10.5539/jel.v7n6p57
- Gaspar, T., Matos, M. G., Ribeiro, J.L., Leal, I., Erhart, M., & Ravens-Sieberer, U. (2012). Health-related quality of life in children and adolescents: subjective well being. *Spanish Journal of Psychology*, 15(1), 177-186. Doi: 10.5209/rev_SJOP.2012.v15.n1.37306

- Golbertein, E., Wen, H., & Miller, B. (2020). Coronavirus Disease 2019 (COVID-19) and Mental Health for Children and Adolescents *JAMA Pediatrics*. doi:10.1001/jamapediatrics.2020.1456
- IREFREA (2020). Consumption of alcohol, tobacco and drugs decreases among young portuguese population during covid19 pandemic. Coimbra: IREFREA Portugal.
- Jensen, D; Drat, J.; Saxena, S.; Arevalo, L.; Matos, M.G.; Kosola, S. (2020 in press) Child and adolescent health needs attention now and in the aftermath of the Covid-19 pandemic. *International Journal of Public Health*.
- Jiménez-Iglesias, A., Camacho, I., Rivera, F., Moreno, C., & Matos, M. (2017). Social Support from Developmental Contexts and Adolescent Substance Use and Well-Being: A Comparative Study of Spain and Portugal. *The Spanish Journal of Psychology*, 20, E64. Doi: 10.1017/sjp.2017.62
- Lewis, A. D., Huebner, E., Malone, P. S., & Valois, R. F. (2011). Life satisfaction and student engagement in adolescents. *Journal of Youth and Adolescence*, 40, 249–262. Doi: 10.1007/s10964-010-9517-6.
- Loureiro, N., & Matos, M. G. (2014). Who are the girls that don't quit: features and particularities of the physically active portuguese female adolescent. *Revista Saúde e Desenvolvimento Humano*, 2(2), 25-38.
- Marques, A., Branquinho, C. & Matos, M. G. (2017). The taller the better? Psychobiological influences on bullying behaviour among Portuguese adolescents. *International Journal of Pediatrics and Neonatal Health*, 1(3), 65-73. doi: 10.25141/2572-4355-2017-3.0065
- Marques, A., Calmeiro, L., Loureiro, L., Frasilho, D., & Matos, M. G. (2015). Health complaints among adolescents: Associations with more screen-based behaviours and less physical activity. *Journal of Adolescence*, 44, 150–157. Doi: 10.1016/j.adolescence.2015.07.018
- Marques, A., Peralta, M., Martins, J., Catunda, R., Matos, M. G., & Saboga Nunes, L. (2016). Associations between physical activity and self-rated wellbeing in European adults: A population-based, cross-sectional study. *Preventive Medicine*, 91, 18-23.
- Marques, S. C., Lopez, S. J., Fontaine, A. M., Coimbra, S., & Mitchell, J. (2015). How much hope is enough? Levels of hope and students' psychological and school functioning. *Psychology in the Schools*, 52, 325–334. Doi: 10.1002/pits.21833.
- Matos, M.G., Tomé, G., Branquinho, C., Reis, M., Ramiro, L., Gomez-Baya, D., Gaspar, T. (2020 in press). *Being Positive, Participative and Flexible: tools to human development and progress*. Erebea.
- Matos, M. G., & Aventura Social Team (2018). A saúde dos adolescentes portugueses após a recessão. Estudo 2018 do Health Behaviour in School Aged Children (HBSC). http://aventurasocial.com/publicacoes/publicacao_1545534554.pdf (ebook).
- Matos, M. G., Camacho, I., Reis, M., Tomé, G., Branquinho, C. & Ramiro, L. (2017). Is truth in the eyes of the beholder? Or are Portuguese schools, as viewed by Portuguese pupils, mismatching with what the educational system offers? *Vulnerable Children and Youth Studies*, 1-11. Doi: 10.1080/17450128.2017.1363447
- Matos, M. G., Marques, A., Calmeiro, L., & Loureiro, N. (2014). Diferentes perfis comportamentais em adolescentes e associação à prática de atividade física [Different behavioral profiles in adolescents and association with physical activity.]. *Psicologia, Saúde & Doença*, 15(2), 495-509. Doi: 10.15309/14psd150213
- Matos, M. G., Reis, M., Camacho, I., Simões, C., Gómez-Baya, D., Mota, C. ... Equipa Aventura Social (2015). Em tempo de recessão, os adolescentes portugueses continuam saudáveis e ou são ainda saudáveis mas já não são felizes? []. In times of recession, Portuguese teenagers are still healthy and or are they still healthy but not happy anymore?]. *Arquivos de Medicina*, 116-122.
- Matos, M. G., Simões, Camacho, Reis, & Aventura Social Team (2015). A saúde dos adolescentes portugueses em tempos de

recessão. Estudo 2014 do Health Behaviour in School Aged Children (HBSC) [The health of Portuguese teenagers in times of recession. 2014 Study of Health Behavior in School Aged Children (HBSC)]. Lisboa: FMH/MS/IHMT.

- Matos, M., Simões, C., Tomé, G., Gaspar, T., Camacho, I., Diniz, J., & Equipa Aventura Social (2006). A saúde dos adolescentes portugueses – Hoje em 8 anos: Relatório Preliminar do estudo HBSC 2006 [The health of Portuguese adolescents - Today in 8 years: Preliminary Report of the HBSC 2006 study]. Retrieved from www.fmh.utl.pt/aventurasocial.com
- Matos, M. G., Simões, C., Tomé, G., Camacho, I., Ferreira, M., Ramiro, L., Reis, M., Gaspar, T., Veloso, S., Loureiro, N., Borges, A., Diniz, J., & Equipa Aventura Social (2012). Aventura Social & Saúde, A Saúde dos adolescentes portugueses – Relatório Final do Estudo HBSC 2010 [Social Adventure & Health, The Health of Portuguese Adolescents - Final Report of the HBSC Study 2010]. Lisboa: Centro Malária e Outras Doenças Tropicais/IHMT/UNL; FMH/Universidade Técnica de Lisboa.
- Roberts, C., Currie, C., Samdal, O., Currie, D., Smith, R., & Maes, L. (2007). Measuring the health behaviours of adolescents through cross-national survey research: recent developments in the Health Behaviour in School-aged Children (HBSC) study. *Journal of Public Health*, 15, 179-186. Doi:10.1007/s10389-007-0100-x.
- Simões, C., Rivera, F., Moreno, C., & Matos, M. (2018). School Performance Paths: Personal and Contextual Factors Related to Top Performers and Low Achievers in Portugal and Spain. *The Spanish Journal of Psychology*, 21, E36. Doi: 10.1017/sjp.2018.37
- Tomé, G., Almeida, A., Ramiro, L., Gaspar, T., & Matos, M.G. (2020, in press). Intervention in Schools promoting mental health and well-being: a systematic review. *Global Journal of Community Psychology Practice*.
- Tomé, G., Matos, M.G., Camacho I., Gomes, P., Reis, M. & Branquinho, C. (2018). Mental Health Promotion in School Context – Validation of the Es'cool scale for teachers. *Journal of Psychiatry and Behavioral Sciences*, 2, 1-11.

